

MAKING NON-TRIVIALY ASSOCIATED MODULAR CATEGORIES FROM FINITE GROUPS

M. M. Al-Shomrani & E. J. Beggs

Department of Mathematics
University of Wales, Swansea
Singleton Park, Swansea SA2 8PP, UK

February 2003

ABSTRACT : We show that the double $\mathcal{D}$ of the non-trivially associated tensor category constructed from left coset representatives of a subgroup of a finite group X is a modular category. Also we give a definition of the character of an object in this category as an element of a braided Hopf algebra in the category. This definition is shown to be adjoint invariant and multiplicative on tensor products. A detailed example is given. Finally we show an equivalence of categories between the non-trivially associated double $\mathcal{D}$ and the trivially associated category of representations of the Drinfeld double of the group $D(X)$.

1 Introduction

This paper will make continual use of formulae and ideas from [2], and these definitions and formulae will not be repeated, as they would add very considerably to the length of the paper. The paper [2] is itself based on the papers [3, 4], but is mostly self contained

in terms of notation and definitions. The book [9] has been used as a standard reference for Hopf algebras, and [13, 1] as references for modular categories.

In [2] there is a construction of a non-trivially associated tensor category $\mathcal{C}$ from data which is a choice of left coset representatives M for a subgroup G of a finite group X . This introduces a binary operation $\cdot$ and a G -valued ‘cocycle’ τ on M . There is also a double construction where X is viewed as a subgroup of a larger group. This gives rise to a braided category $\mathcal{D}$, which is the category of reps of an algebra D , which is itself in the category, and it is the category that we concentrate on in this paper.

It is our aim to show that the non-trivially associated algebra D has reps which have characters in the same way that the reps of a finite group have characters, and also that the category of its representations has a modular structure in the same way that the category of reps of the double of a group has a modular structure.

We begin by describing the indecomposable objects in $\mathcal{C}$, in a similar manner to that used in [4]. A detailed example is given using the group D_6 . Then we show how to find the dual objects in the category, and again illustrate this with the example.

Next we show that the rigid braided category $\mathcal{D}$ is a ribbon category. The ribbon maps are calculated for the indecomposable objects in our example category.

In the next section we explicitly evaluate in $\mathcal{D}$ the standard diagram for trace in a ribbon category [9]. Then we define the character of an object in $\mathcal{D}$ as an element of the dual of the braided Hopf algebra D . This element is shown to be right adjoint invariant. Also we show that the character is multiplicative for the tensor product of objects. A formula is found for the character in $\mathcal{D}$ in terms of characters of group representations.

The last ingredient needed for a modular category is the trace of the double braiding,

and this is calculated in $\mathcal{D}$ in terms of group characters. Then the matrices S , T and C implementing the modular representation are calculated explicitly in our example.

Finally we show an equivalence of categories between the non-trivially associated double $\mathcal{D}$ and the category of representations of the Drinfeld double of the group $D(X)$.

Throughout the paper we assume that all groups mentioned are finite, and that all vector spaces are finite dimensional. We take the base field to be the complex numbers $\mathbb{C}$.

2 Indecomposable objects in $\mathcal{C}$

The objects of $\mathcal{C}$ are the right representations of the algebra A described in [2]. We now look at the indecomposable objects in $\mathcal{C}$, or the irreducible representations of A , in a manner similar to that used in [4].

Theorem 2.1 *The indecomposable objects in $\mathcal{C}$ are of the form*

$$V = \bigoplus_{s \in \mathcal{O}} V_s$$

where $\mathcal{O}$ is an orbit in M under the G action $\triangleleft$, and each V_s is an irreducible right representation of the stabilizer of s , $\text{stab}(s)$. Every object T in $\mathcal{C}$ can be written as a direct sum of indecomposable objects in $\mathcal{C}$.

Proof. For an object T in $\mathcal{C}$ we can use the M -grading to write

$$T = \bigoplus_{s \in M} T_s, \tag{1}$$

but as M is a disjoint union of orbits $\mathcal{O}_s = \{s \triangleleft u : u \in G\}$ for $s \in M$, T can be rewritten as a disjoint sum over orbits,

$$T = \bigoplus_{\mathcal{O}} T_{\mathcal{O}}, \tag{2}$$

where

$$T_{\mathcal{O}} = \bigoplus_{s \in \mathcal{O}} T_s. \quad (3)$$

Now we will define the stabilizer of $s \in \mathcal{O}$, which is a subgroup of G , as

$$\text{stab}(s) = \{u \in G : s \triangleleft u = s\}.$$

As $\langle \eta \bar{\triangleleft} u \rangle = \langle \eta \rangle \triangleleft u$ for all $\eta \in T$, T_s is a representation of the group $\text{stab}(s)$. Now fix a base point $t \in \mathcal{O}$. Because $\text{stab}(t)$ is a finite group, T_t is a direct sum of irreducible group representations W_i for $i = 1, \dots, m$, i.e.,

$$T_t = \bigoplus_{i=1}^m W_i. \quad (4)$$

Suppose that $\mathcal{O} = \{t_1, t_2, \dots, t_n\}$ where $t_1 = t$, and take $u_i \in G$ so that $t_i = t \triangleleft u_i$. Define

$$U_i = \bigoplus_{j=1}^n W_i \bar{\triangleleft} u_j \subset \bigoplus_{s \in \mathcal{O}} T_s. \quad (5)$$

We claim that each U_i is an indecomposable object in $\mathcal{C}$. For any $v \in G$ and $\xi \bar{\triangleleft} u_k \in W_i \bar{\triangleleft} u_k$,

$$(\xi \bar{\triangleleft} u_k) \bar{\triangleleft} v = (\xi \bar{\triangleleft} (u_k v u_j^{-1})) \bar{\triangleleft} u_j,$$

where $u_k v u_j^{-1} \in \text{stab}(t)$ for some $u_j \in G$. This shows that U_i is a representation of G .

By the definition of U_i , any subrepresentation of U_i which contains W_i must be all of U_i .

Thus U_i is an indecomposable object in $\mathcal{C}$, and

$$T_{\mathcal{O}} = \bigoplus_{i=1}^m U_i. \quad \square \quad (6)$$

Theorem 2.2 {Schur's lemma} *Let V and W be two indecomposable objects in $\mathcal{C}$, and let $\alpha : V \longrightarrow W$ be a morphism. Then α is zero or a scalar multiple of the identity.*

Proof. V and W are associated to orbits $\mathcal{O}$ and $\mathcal{O}'$ so that $V = \bigoplus_{s \in \mathcal{O}} V_s$ and $W = \bigoplus_{s \in \mathcal{O}'} W_s$. As morphisms preserve grade, if $\alpha \neq 0$, then $\mathcal{O} = \mathcal{O}'$. Now if we take $s \in \mathcal{O}$, we find that $\alpha : V_s \rightarrow W_s$ is a map of irreps of $\text{stab}(s)$, so by Schur's lemma for groups, any non-zero map is a scalar multiple of the identity, and we have $V_s = W_s$ as representations of $\text{stab}(s)$. Now we need to check that the multiple of the identity is the same for each $s \in \mathcal{O}$. Suppose α is a multiplication by λ on V_s . Given $t \in \mathcal{O}$, there is a $u \in G$ so that $t \triangleleft u = s$. Then for $\eta \in V_t$,

$$\alpha(\eta) = \alpha(\eta \bar{\triangleleft} u) \bar{\triangleleft} u^{-1} = \lambda(\eta \bar{\triangleleft} u) \bar{\triangleleft} u^{-1} = \lambda \eta. \quad \square$$

Lemma 2.3 *Let V be an indecomposable object in $\mathcal{C}$ associated to the orbit $\mathcal{O}$. Choose $s, t \in \mathcal{O}$ and $u \in G$ so that $s \triangleleft u = t$. Then V_s and V_t are irreps of $\text{stab}(s)$ and $\text{stab}(t)$ respectively, and the group characters obey $\chi_{V_t}(v) = \chi_{V_s}(u v u^{-1})$.*

Proof. Note that $\bar{\triangleleft} u$ is an invertible map from V_s to V_t . Then we have the commuting diagram

$$\begin{array}{ccc} V_s & \xrightarrow{\bar{\triangleleft} u v u^{-1}} & V_s \\ \downarrow \bar{\triangleleft} u & & \downarrow \bar{\triangleleft} u \\ V_t & \xrightarrow{\bar{\triangleleft} v} & V_t \end{array}$$

which implies that $\text{trace}(\bar{\triangleleft} u v u^{-1} : V_s \rightarrow V_s) = \text{trace}(\bar{\triangleleft} v : V_t \rightarrow V_t)$. $\square$

3 An example of indecomposable objects

We give an example of indecomposable objects in the categories discussed in the last section. As we will later want to have a category with braiding, we use the double construction in [2]. We also use lemma 2.3 to list the group characters [5] for every point in the orbit in terms of the given base points.

Take X to be the dihedral group $D_6 = \langle a, b : a^6 = b^2 = e, ab = ba^5 \rangle$, whose elements we list as $\{e, a, a^2, a^3, a^4, a^5, b, ba, ba^2, ba^3, ba^4, ba^5\}$, and G to be the non-abelian normal subgroup of order 6 generated by a^2 and b , i.e. $G = \{e, a^2, a^4, b, ba^2, ba^4\}$. We choose $M = \{e, a\}$. The center of D_6 is the subgroup $\{e, a^3\}$, and it has the following conjugacy classes: $\{e\}$, $\{a^3\}$, $\{a^2, a^4\}$, $\{a, a^5\}$, $\{b, ba^2, ba^4\}$ and $\{ba, ba^3, ba^5\}$.

The category $\mathcal{D}$ consists of right representations of the group $X = D_6$ which are graded by $Y = D_6$ (as a set), using the actions $\tilde{\alpha} : Y \times X \rightarrow Y$ and $\tilde{\beta} : Y \times X \rightarrow X$ which are defined as follows:

$$y\tilde{\alpha}x = x^{-1}yx, \quad \text{and} \quad vt\tilde{\beta}x = v^{-1}xv' = txt'^{-1},$$

for $x \in X$, $y \in Y$, $v, v' \in G$ and $t, t' \in M$ where $vt\tilde{\alpha}x = v't'$.

Now let V be an indecomposable object in $\mathcal{D}$. We get the following cases:

Case (1): Take the orbit $\{e\}$ with base point e , whose stabilizer is the whole of D_6 . There are six possible irreducible group representations of the stabilizer, with their characters given by table(1) [12]:

irreps		$\{e\}$	$\{a^3\}$	$\{b, ba^2, ba^4\}$	$\{ba, ba^3, ba^5\}$	$\{a^2, a^4\}$	$\{a, a^5\}$
1_1	2_1	1	1	1	1	1	1
1_2	2_2	1	-1	-1	1	1	-1
1_3	2_3	1	-1	1	-1	1	-1
1_4	2_4	1	1	-1	-1	1	1
1_5	2_5	2	-2	0	0	-1	1
1_6	2_6	2	2	0	0	-1	-1

table (1)

Case (2): Take the orbit $\{a^3\}$ with base point a^3 , whose stabilizer is the whole of D_6 . There are six possible irreps $\{2_1, 2_2, 2_3, 2_4, 2_5, 2_6\}$, with characters given by table(1).

Case (3): Take the orbit $\{a^2, a^4\}$ with base point a^2 , whose stabilizer is $\{e, a, a^2, a^3, a^4, a^5\}$. There are six irreps $\{3_0, 3_1, 3_2, 3_3, 3_4, 3_5\}$, with characters given by table(2), where

$\omega = e^{i\pi/3}$. Applying lemma 2.3 gives $\chi_{V_{a^4}}(v) = \chi_{V_{a^2}}(bvb)$.

irreps		e	a	a^2	a^3	a^4	a^5
3_0	4_0	1	1	1	1	1	1
3_1	4_1	1	ω^1	ω^2	ω^3	ω^4	ω^5
3_2	4_2	1	ω^2	ω^4	1	ω^2	ω^4
3_3	4_3	1	ω^3	1	ω^3	1	ω^3
3_4	4_4	1	ω^4	ω^2	1	ω^4	ω^2
3_5	4_5	1	ω^5	ω^4	ω^3	ω^2	ω^1

table (2)

Case (4): Take the orbit $\{a, a^5\}$ with base point a , whose stabilizer is $\{e, a, a^2, a^3, a^4, a^5\}$.

There are six irreps $\{4_0, 4_1, 4_2, 4_3, 4_4, 4_5\}$ with characters given in table(2). Applying lemma 2.3 gives $\chi_{V_{a^5}}(v) = \chi_{V_a}(ba^2vba^2)$.

Case (5): Take the orbit $\{b, ba^2, ba^4\}$ with base point b , whose stabilizer is $\{e, a^3, b, ba^3\}$.

There are four irreps with characters given by table(3). Applying lemma 2.3 gives $\chi_{V_{ba^2}}(v) = \chi_{V_b}(a^4va^2)$ and $\chi_{V_{ba^4}}(v) = \chi_{V_b}(a^2va^4)$.

	e	a^3	b	ba^3
5_{++}	1	1	1	1
5_{+-}	1	1	-1	-1
5_{-+}	1	-1	1	-1
5_{--}	1	-1	-1	1

table (3)

Case (6): Take the orbit $\{ba, ba^3, ba^5\}$ with base point ba , whose stabilizer is $\{e, a^3, ba, ba^4\}$.

There are four irreps with character given by table(4). Applying lemma 2.3 gives $\chi_{V_{ba^3}}(v) = \chi_{V_{ba}}(a^4va^2)$ and $\chi_{V_{ba^5}}(v) = \chi_{V_{ba}}(a^2va^4)$.

	e	a^3	ba	ba^4
6_{++}	1	1	1	1
6_{-+}	1	-1	1	-1
6_{+-}	1	1	-1	-1
6_{--}	1	-1	-1	1

table (4)

4 Duals of indecomposable objects in $\mathcal{C}$

Given an irreducible object V with associated orbit $\mathcal{O}$ in $\mathcal{C}$, how do we find its dual V^* ? The dual would be described as in section 2, by an orbit, a base point in the orbit and a right group representation of the stabilizer of the base point. Using the formula $(s^L \cdot s) \triangleleft u = (s^L \triangleleft (s \triangleright u)) \cdot (s \triangleleft u) = e$, we see that the left inverse of a point in the orbit containing s is in the orbit containing s^L . By using the evaluation map from $V^* \otimes V$ to the field we can take $(V^*)_{s^L} = (V_s)^*$ as vector spaces. We use $\check{\triangleleft}$ as the action of $\text{stab}(s)$ on $(V_s)^*$, i.e. $(\alpha \check{\triangleleft} z)(\xi \bar{\triangleleft} z) = \alpha(\xi)$ for $\alpha \in (V_s)^*$ and $\xi \in V_s$. The action $\bar{\triangleleft}$ of $\text{stab}(s^L)$ on $(V^*)_{s^L}$ is given by $\alpha \bar{\triangleleft} (s \triangleright z) = \alpha \check{\triangleleft} z$ for $z \in \text{stab}(s)$. In terms of group characters this gives

$$\chi_{(V^*)_{s^L}}(s \triangleright z) = \chi_{(V_s)^*}(z), \quad z \in \text{stab}(s).$$

If we take $\mathcal{O}^L = \{s^L : s \in \mathcal{O}\}$ to have base point p , and choose $u \in G$ so that $p \triangleleft u = s^L$, then using lemma 2.3 gives

$$\chi_{(V^*)_{s^L}}(s \triangleright z) = \chi_{(V_s)^*}(z) = \chi_{(V^*)_p}(u(s \triangleright z)u^{-1}), \quad z \in \text{stab}(s). \quad (7)$$

This formula allows us to find the character of V^* at its base point p as a representation of $\text{stab}(p)$ in terms of the character of the dual of V_s as a representation of $\text{stab}(s)$.

Lemma 4.1 *In $\mathcal{C}$ we can regard the dual $(V \otimes W)^*$ as $W^* \otimes V^*$ with the evaluation*

$$(\alpha \otimes \beta)(\xi \otimes \eta) = (\alpha \bar{\triangleleft} \tau(\langle \beta \rangle, \langle \xi \rangle \cdot \langle \eta \rangle))(\eta) (\beta \bar{\triangleleft} \tau(\langle \xi \rangle, \langle \eta \rangle)^{-1})(\xi).$$

Given a basis $\{\xi\}$ of V and a basis $\{\eta\}$ of W , the dual basis $\{\widehat{\xi \otimes \eta}\}$ of $W^ \otimes V^*$ can be written in terms of the dual basis of V^* and W^* as*

$$\widehat{\xi \otimes \eta} = \hat{\eta} \bar{\triangleleft} \tau(\langle \xi \rangle^L \triangleleft \tau(\langle \xi \rangle, \langle \eta \rangle), \langle \xi \rangle \cdot \langle \eta \rangle)^{-1} \otimes \hat{\xi} \bar{\triangleleft} \tau(\langle \xi \rangle, \langle \eta \rangle).$$

Proof. Applying the associator to $(\alpha \otimes \beta) \otimes (\xi \otimes \eta)$ gives

$$\alpha \bar{\Delta} \tau(\langle \beta \rangle, \langle \xi \rangle \cdot \langle \eta \rangle) \otimes (\beta \otimes (\xi \otimes \eta)) ,$$

and then applying the inverse associator gives

$$\alpha \bar{\Delta} \tau(\langle \beta \rangle, \langle \xi \rangle \cdot \langle \eta \rangle) \otimes \left((\beta \bar{\Delta} \tau(\langle \xi \rangle, \langle \eta \rangle)^{-1} \otimes \xi) \otimes \eta \right) .$$

Applying the evaluation map first to $\beta \bar{\Delta} \tau(\langle \xi \rangle, \langle \eta \rangle)^{-1} \otimes \xi$, then to $\alpha \bar{\Delta} \tau(\langle \beta \rangle, \langle \xi \rangle \cdot \langle \eta \rangle) \otimes \eta$ gives the first equation. For the evaluation to be non-zero, we need $(\langle \beta \rangle \triangleleft \tau(\langle \xi \rangle, \langle \eta \rangle)^{-1}) \cdot \langle \xi \rangle = e$ which implies $\langle \beta \rangle \triangleleft \tau(\langle \xi \rangle, \langle \eta \rangle)^{-1} = \langle \xi \rangle^L$, or equivalently $\langle \beta \rangle = \langle \xi \rangle^L \triangleleft \tau(\langle \xi \rangle, \langle \eta \rangle)$. This gives the second equation. $\square$

Example 4.2 Using (7) we calculate the duals of the objects given in the last section.

Case (1): The orbit $\{e\}$ has left inverse $\{e\}$, so $\chi_{(V^*)_e} = \chi_{(V_e)^*}$. By a calculation with group characters, all the listed irreps of $\text{stab}(e)$ are self-dual, so $1_r^* = 1_r$ for $r \in \{1, \dots, 6\}$.

Case (2): The orbit $\{a^3\}$ has left inverse $\{a^3\}$, so $\chi_{(V^*)_{a^3}} = \chi_{(V_{a^3})^*}$. As in the last case the group representations are self-dual, so $2_r^* = 2_r$ for $r \in \{1, \dots, 6\}$.

Case (3): The left inverse of the base point a^2 is a^4 , which is still in the orbit. As group representations, the dual of 3_r is $3_{6-r} \pmod{6}$. Applying lemma 2.3 to move the base point, we see that the dual of 3_r in the category is 3_r .

Case (4): The left inverse of the base point a is a^5 , which is still in the orbit. As in the last case, the dual of 4_r in the category is 4_r .

Case (5): The left inverse of the base point is itself, and as group representations, all case 5 irreps are self dual. We deduce that in the category the objects are self dual.

Case (6): Self dual, as in case 5.

5 The ribbon map on the category $\mathcal{D}$

Theorem 5.1 *The ribbon transformation $\theta_V : V \longrightarrow V$ for any object V in $\mathcal{D}$ can be defined by $\theta_V(\xi) = \xi \hat{\triangleleft} \|\xi\|$.*

Proof. In the following lemmas we show that the required properties hold. $\square$

Lemma 5.2 *θ_V is a morphism in the category.*

Proof. Begin by checking the X -grade, for $\xi \in V$

$$\|\theta_V(\xi)\| = \|\xi \hat{\triangleleft} \|\xi\|\| = \|\xi\| \hat{\triangleleft} \|\xi\| = \|\xi\|.$$

Now we check the X -action, i.e. that $\theta_V(\xi \hat{\triangleleft} x) = \theta_V(\xi) \hat{\triangleleft} x$.

$$\begin{aligned} \theta_V(\xi \hat{\triangleleft} x) &= (\xi \hat{\triangleleft} x) \hat{\triangleleft} \|\xi \hat{\triangleleft} x\| = (\xi \hat{\triangleleft} x) \hat{\triangleleft} (\|\xi\| \hat{\triangleleft} x) \\ &= \xi \hat{\triangleleft} x x^{-1} \|\xi\| x = (\xi \hat{\triangleleft} \|\xi\|) \hat{\triangleleft} x = \theta_V(\xi) \hat{\triangleleft} x. \end{aligned} \quad \square$$

Lemma 5.3 *For any two objects V and W in $\mathcal{D}$,*

$$\theta_{V \otimes W} = \Psi_{V \otimes W}^{-1} \circ \Psi_{W \otimes V}^{-1} \circ (\theta_V \otimes \theta_W) = (\theta_V \otimes \theta_W) \circ \Psi_{V \otimes W}^{-1} \circ \Psi_{W \otimes V}^{-1}$$

This can also be described by figure 1:




figure 1

Proof. First calculate $\Psi(\Psi(\xi \otimes \eta))$ for $\xi \in V$ and $\eta \in W$, beginning with

$$\Psi(\Psi(\xi \otimes \eta)) = \Psi(\eta \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\eta|)^{-1} \otimes \xi \hat{\triangleleft} |\eta|). \quad (8)$$

To simplify what follows we shall use the substitutions

$$\eta' = \xi \hat{\triangleleft} |\eta| \quad \text{and} \quad \xi' = \eta \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\eta|)^{-1}, \quad (9)$$

so equation (8) can be rewritten as

$$\begin{aligned} \Psi(\Psi(\xi \otimes \eta)) &= \Psi(\xi' \otimes \eta') \\ &= \eta' \hat{\triangleleft} (\langle \xi' \rangle \triangleleft |\eta'|)^{-1} \otimes \xi' \hat{\triangleleft} |\eta'|. \end{aligned} \quad (10)$$

As $\eta' = \xi \hat{\triangleleft} |\eta| = \xi \triangleleft |\eta|$, then $|\eta'| = |\xi \triangleleft |\eta|| = (\langle \xi \rangle \triangleright |\eta|)^{-1} |\xi| |\eta|$, so

$$\begin{aligned} \xi' \hat{\triangleleft} |\eta'| &= \eta \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\eta|)^{-1} (\langle \xi \rangle \triangleright |\eta|)^{-1} |\xi| |\eta| \\ &= \eta \hat{\triangleleft} ((\langle \xi \rangle \triangleright |\eta|)(\langle \xi \rangle \triangleleft |\eta|))^{-1} |\xi| |\eta| \\ &= \eta \hat{\triangleleft} |\eta|^{-1} \langle \xi \rangle^{-1} |\xi| |\eta|. \end{aligned} \quad (11)$$

Hence if we put $y = \|\xi \otimes \eta\| = \|\xi\| \circ \|\eta\| = |\eta|^{-1} |\xi|^{-1} \langle \xi \rangle \langle \eta \rangle$,

$$\Psi(\Psi(\xi \otimes \eta)) \hat{\triangleleft} \|\xi \otimes \eta\| = \xi \hat{\triangleleft} |\eta| (\langle \xi' \rangle \triangleleft |\eta'|)^{-1} (p \tilde{\triangleright} \|\xi \otimes \eta\|) \otimes \eta \hat{\triangleleft} |\eta|^{-1} \langle \eta \rangle, \quad (12)$$

where, using (11),

$$p = \|\xi' \hat{\triangleleft} |\eta'|\| = |\xi' \triangleleft |\eta'||^{-1} \langle \xi' \rangle \triangleleft |\eta'| = \|\eta\| \tilde{\triangleleft} \|\eta\| y^{-1} = \|\eta\| \tilde{\triangleleft} y^{-1}$$

$$p \tilde{\triangleright} \|\xi \otimes \eta\| = (\|\eta\| \tilde{\triangleleft} y^{-1}) \tilde{\triangleright} y = (\|\eta\| \tilde{\triangleright} y^{-1})^{-1},$$

As $\|\xi' \triangleleft |\eta'|\| = v' t' = \|\eta\| \tilde{\triangleleft} y^{-1}$, by unique factorization, $t' = \langle \xi' \rangle \triangleleft |\eta'|$. Then $\|\eta\| \tilde{\triangleright} y^{-1} = \langle \eta \rangle y^{-1} t'^{-1}$, which implies that

$$|\eta| (\langle \xi' \rangle \triangleleft |\eta'|)^{-1} (\|\eta\| \tilde{\triangleright} y^{-1})^{-1} = |\eta| t'^{-1} t' y \langle \eta \rangle^{-1} = \|\xi\|. \quad (13)$$

Substituting this in (12) gives

$$\Psi(\Psi(\xi \otimes \eta)) \hat{\triangleleft} \|\xi \otimes \eta\| = \xi \hat{\triangleleft} \|\xi\| \otimes \eta \hat{\triangleleft} \|\eta\|. \quad \square$$

Lemma 5.4 *For the unit object $\underline{1} = \mathbb{C}$ in $\mathcal{D}$, $\theta_{\underline{1}}$ is the identity.*

Proof. For any object V in $\mathcal{D}$, $\theta_V : V \longrightarrow V$ is defined by

$$\theta_V(\xi) = \xi \hat{\triangleleft} \|\xi\| \quad \text{for } \xi \in V.$$

If we choose $V = \underline{1} = \mathbb{C}$ then $\theta_{\underline{1}}(\xi) = \xi \hat{\triangleleft} e = \xi$ as $\|\xi\| = e$. $\square$

Lemma 5.5 *For any object V in $\mathcal{D}$, $(\theta_V)^* = \theta_{V^*}$ (see figure 2).*




figure 2

Proof. Begin with

$$\text{coev}_V(1) = \sum_{\xi \in \text{basis of } V} \xi \hat{\triangleleft} \tilde{\tau}(\|\xi\|^L, \|\xi\|)^{-1} \otimes \hat{\xi} = \sum_{\xi \in \text{basis of } V} \xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \otimes \hat{\xi}$$

For $\alpha \in V^*$, we follow figure 2 and calculate

$$(\theta_V)^*(\alpha) = (\text{eval}_V \otimes id) \sum_{\xi \in \text{basis of } V} \Phi^{-1} \left(\alpha \otimes \left(\theta_V(\xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}) \otimes \hat{\xi} \right) \right). \quad (14)$$

Now as $\tau(\langle \xi \rangle^L, \langle \xi \rangle) = \langle \xi \rangle^L \langle \xi \rangle$,

$$\begin{aligned}
\| \xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \| &= \| \xi \| \hat{\triangleleft} (\langle \xi \rangle^L \langle \xi \rangle)^{-1} \\
&= \langle \xi \rangle^L \langle \xi \rangle |\xi|^{-1} \langle \xi \rangle \langle \xi \rangle^{-1} \langle \xi \rangle^{L-1} \\
&= \langle \xi \rangle^L \langle \xi \rangle |\xi|^{-1} \langle \xi \rangle^{L-1}, \\
\theta_V(\xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}) &= (\xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}) \hat{\triangleleft} \| \xi \hat{\triangleleft} \tilde{\tau}(\| \xi \|^L, \| \xi \|)^{-1} \| \\
&= \xi \hat{\triangleleft} \langle \xi \rangle^{-1} \langle \xi \rangle^{L-1} \langle \xi \rangle^L \langle \xi \rangle |\xi|^{-1} \langle \xi \rangle^{L-1} \\
&= \xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1}.
\end{aligned}$$

The next step is to find

$$\begin{aligned}
\Phi^{-1}\left(\alpha \otimes \left((\xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1}) \otimes \hat{\xi}\right)\right) &= \left(\alpha \hat{\triangleleft} \tilde{\tau}(\| \xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1} \|, \| \hat{\xi} \|)^{-1}\right. \\
&\quad \left. \otimes (\xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1})\right) \otimes \hat{\xi}.
\end{aligned}$$

As

$$\begin{aligned}
\| \xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1} \| &= \| \xi \| \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1} \\
&= \langle \xi \rangle^L |\xi| |\xi|^{-1} \langle \xi \rangle |\xi|^{-1} \langle \xi \rangle^{L-1} \\
&= \tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1} \langle \xi \rangle^{L-1} \\
&= \tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1} \langle \xi \rangle \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \\
&= \tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1} (\langle \xi \rangle \triangleright \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}) (\langle \xi \rangle \triangleleft \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}),
\end{aligned}$$

then as $\| \hat{\xi} \| = \| \xi \|^L = |\xi| \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \langle \xi \rangle^L$,

$$\begin{aligned}
\Phi^{-1}\left(\alpha \otimes \left((\xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1}) \otimes \hat{\xi}\right)\right) &= \left(\alpha \hat{\triangleleft} \tau(\langle \xi \rangle \triangleleft \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}, \langle \xi \rangle^L)^{-1}\right. \\
&\quad \left. \otimes (\xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1})\right) \otimes \hat{\xi}.
\end{aligned}$$

Put $v = \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} = \langle \xi \rangle^{-1} \langle \xi \rangle^{L-1}$ and $w = \tau(\langle \xi \rangle \triangleleft v, \langle \xi \rangle^L)^{-1} = ((\langle \xi \rangle \triangleleft v) \langle \xi \rangle^L)^{-1}$, then

substituting in (14) gives

$$(\theta_V)^*(\alpha) = (\text{eval}_V \otimes \text{id}) \sum_{\xi \in \text{basis of } V} \left((\alpha \hat{\triangleleft} w) \otimes (\xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1}) \right) \otimes \hat{\xi}. \quad (15)$$

For a given term in the sum to be non-zero, we require

$$\|\alpha\| = \|\hat{\xi}\| = \|\xi\|^L = |\xi| \langle \xi \rangle^{-1} , \quad (16)$$

and we proceed under this assumption. Now calculate

$$\text{eval}_V((\alpha \hat{\triangleleft} w) \otimes (\xi \hat{\triangleleft} |\xi|^{-1} \langle \xi \rangle^{L-1})) = (\beta \hat{\triangleleft} (\|\xi\| \tilde{\triangleright} p))(\xi \tilde{\triangleleft} p) = \beta(\xi) \quad (17)$$

where $p = |\xi|^{-1} \langle \xi \rangle^{L-1}$ and $\beta = \alpha \hat{\triangleleft} w (\|\xi\| \tilde{\triangleright} p)^{-1}$. Next we want to find $\|\xi\| \tilde{\triangleright} p$. To do this, we first find

$$\begin{aligned} \|\xi\| \tilde{\triangleleft} p &= \langle \xi \rangle^L |\xi| |\xi|^{-1} \langle \xi \rangle |\xi|^{-1} \langle \xi \rangle^{L-1} \\ &= v^{-1} |\xi|^{-1} \langle \xi \rangle v = v^{-1} |\xi|^{-1} (\langle \xi \rangle \triangleright v) (\langle \xi \rangle \triangleleft v) , \end{aligned} \quad (18)$$

and hence

$$\begin{aligned} \|\xi\| \tilde{\triangleright} p &= \langle \xi \rangle p (\langle \xi \rangle \triangleleft v)^{-1} \\ &= \langle \xi \rangle |\xi|^{-1} \langle \xi \rangle v (\langle \xi \rangle \triangleleft v)^{-1} \\ &= \langle \xi \rangle |\xi|^{-1} (\langle \xi \rangle \triangleright v) . \end{aligned} \quad (19)$$

Thus

$$\begin{aligned} \beta &= \alpha \hat{\triangleleft} w (\langle \xi \rangle \triangleright v)^{-1} |\xi| \langle \xi \rangle^{-1} \\ &= \alpha \hat{\triangleleft} \langle \xi \rangle^{L-1} (\langle \xi \rangle \triangleleft v)^{-1} (\langle \xi \rangle \triangleright v)^{-1} |\xi| \langle \xi \rangle^{-1} \\ &= \alpha \hat{\triangleleft} \langle \xi \rangle v (\langle \xi \rangle v)^{-1} |\xi| \langle \xi \rangle^{-1} = \alpha \hat{\triangleleft} |\xi| \langle \xi \rangle^{-1} . \end{aligned} \quad (20)$$

Now substituting these last equations in (15) gives

$$(\theta_V)^*(\alpha) = \sum_{\xi \in \text{basis of } V \text{ with } |\xi| \langle \xi \rangle^{-1} = \|\alpha\|} (\alpha \hat{\triangleleft} \|\alpha\|)(\xi) \cdot \hat{\xi} \quad (21)$$

Take a basis $\xi_1, \xi_2, \dots, \xi_n$ with $(\alpha \hat{\triangleleft} \|\alpha\|)(\xi_i)$ being 1 if $i = 1$, and 0 otherwise. Then

$$(\theta_V)^*(\alpha) = \hat{\xi}_1 + 0 = \alpha \hat{\triangleleft} \|\alpha\| = \theta_{V^*}(\alpha) ,$$

where $\hat{\xi}_1, \hat{\xi}_2, \dots, \hat{\xi}_n$ is the dual basis of V^* defined by $\hat{\xi}_i(\xi_j) = \delta_{i,j}$. $\square$

Example 5.6 We return to the example of section 3. First we calculate the value of the ribbon map on the indecomposable objects. For an irreducible representation V , we have $\theta_V : V \rightarrow V$ defined by $\theta_V(\xi) = \xi \hat{\Delta} \|\xi\|$ for $\xi \in V$. At the base point $s \in \mathcal{O}$, we have $\theta_V(\xi) = \xi \bar{\Delta} s$ for $\xi \in V$ and $\theta : V_s \rightarrow V_s$ is a multiple Θ_V , say, of the identity or, in more explicitly, $\text{trace}(\theta : V_s \rightarrow V_s) = \Theta_V \dim_{\mathbb{C}}(V_s)$, i.e.,

$$\Theta_V = \frac{\text{group character}(s)}{\dim_{\mathbb{C}}(V_s)} \quad (22)$$


And then, for the different cases we will get the following table :

irreps	Θ_V	irreps	Θ_V
1_1	1	3_4	ω^2
1_2	1	3_5	ω^4
1_3	1	4_0	1
1_4	1	4_1	ω^1
1_5	1	4_2	ω^2
1_6	1	4_3	-1
2_1	1	4_4	ω^4
2_2	-1	4_5	ω^5
2_3	-1	5_{++}	1
2_4	1	5_{+-}	-1
2_5	-1	5_{-+}	1
2_6	1	5_{--}	-1
3_0	1	6_{++}	1
3_1	ω^2	6_{-+}	1
3_2	ω^4	6_{+-}	-1
3_3	1	6_{--}	-1

table (5)

6 Traces in the category $\mathcal{D}$

Definition 6.1 [13] The trace of a morphism $T : V \longrightarrow V$ for any object V in $\mathcal{D}$ is defined by



Theorem 6.2 *If we evaluate the diagram of definition 6.1 in $\mathcal{D}$, we find*

$$\text{trace}(T) = \sum_{\xi \in \text{basis of } V} \hat{\xi} (T(\xi)) .$$

Proof. Begin with

$$\text{coev}_V(1) = \sum_{\xi \in \text{basis of } V} \xi \hat{\Delta} \tilde{\tau}(\|\xi\|^L, \|\xi\|)^{-1} \otimes \hat{\xi} = \sum_{\xi \in \text{basis of } V} \xi \hat{\Delta} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \otimes \hat{\xi},$$

and applying $T \otimes \text{id}$ to this gives

$$\sum_{\xi \in \text{basis of } V} T(\xi \hat{\Delta} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}) \otimes \hat{\xi} = \sum_{\xi \in \text{basis of } V} T(\xi) \hat{\Delta} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \otimes \hat{\xi}.$$

Next apply the braiding map to the last equation to get

$$\sum_{\xi \in \text{basis of } V} \Psi(T(\xi) \hat{\Delta} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \otimes \hat{\xi}) = \sum_{\xi \in \text{basis of } V} \hat{\xi} \hat{\Delta} (\langle \xi' \rangle \triangleleft |\hat{\xi}|)^{-1} \otimes \xi' \hat{\Delta} |\hat{\xi}| \quad (23)$$

where $\xi' = T(\xi) \hat{\Delta} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}$, so

$$\begin{aligned} \langle \xi' \rangle &= \langle T(\xi) \hat{\Delta} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \rangle = \langle T(\xi) \bar{\Delta} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \rangle \\ &= \langle T(\xi) \rangle \triangleleft \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} = \langle \xi \rangle \triangleleft \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}. \end{aligned} \quad (24)$$

To calculate $|\hat{\xi}|$ we start with

$$\|\hat{\xi}\| = \|\xi\|^L = (|\xi|^{-1} \langle \xi \rangle)^L = |\xi| \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \langle \xi \rangle^L,$$

which implies that $|\hat{\xi}| = \tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1}$. Then

$$\begin{aligned} \hat{\xi} \hat{\triangleleft} (\langle \xi' \rangle \triangleleft |\hat{\xi}|)^{-1} &= \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1})^{-1} \\ &= \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1}, \\ \xi' \triangleleft |\hat{\xi}| &= (T(\xi) \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}) \hat{\triangleleft} (\tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1}) = T(\xi) \hat{\triangleleft} |\xi|^{-1}, \end{aligned}$$

which gives

$$\sum_{\xi \in \text{basis of } V} \hat{\xi} \hat{\triangleleft} (\langle \xi' \rangle \triangleleft |\hat{\xi}|)^{-1} \otimes \xi' \hat{\triangleleft} |\hat{\xi}| = \sum_{\xi \in \text{basis of } V} \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes T(\xi) \hat{\triangleleft} |\xi|^{-1}. \quad (25)$$

Next

$$\begin{aligned} \theta^{-1}(T(\xi) \hat{\triangleleft} |\xi|^{-1}) &= (T(\xi) \hat{\triangleleft} |\xi|^{-1}) \hat{\triangleleft} \|T(\xi) \hat{\triangleleft} |\xi|^{-1}\|^{-1} \\ &= (T(\xi) \hat{\triangleleft} |\xi|^{-1}) \hat{\triangleleft} (\|T(\xi)\| \tilde{\triangleleft} |\xi|^{-1})^{-1} \\ &= T(\xi) \hat{\triangleleft} |\xi|^{-1} (\|\xi\| \tilde{\triangleleft} |\xi|^{-1})^{-1} \\ &= T(\xi) \hat{\triangleleft} |\xi|^{-1} (|\xi| |\xi|^{-1} \langle \xi \rangle |\xi|^{-1})^{-1} \\ &= T(\xi) \hat{\triangleleft} |\xi|^{-1} |\xi| \langle \xi \rangle^{-1} = T(\xi) \hat{\triangleleft} \langle \xi \rangle^{-1}, \end{aligned} \quad (26)$$

and finally we need to calculate

$$\text{eval}(\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes T(\xi) \hat{\triangleleft} \langle \xi \rangle^{-1}) = (\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1}) (T(\xi) \hat{\triangleleft} \langle \xi \rangle^{-1}). \quad (27)$$

We know, from the definition of the action on V^* , that

$$(\hat{\xi} \hat{\triangleleft} (\|T(\xi)\| \tilde{\triangleright} x)) (T(\xi) \hat{\triangleleft} x) = \hat{\xi}(T(\xi)). \quad (28)$$

If we put $x = \langle \xi \rangle^{-1}$, we want to show that $\|T(\xi)\| \tilde{\triangleright} x = (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1}$, so

$$\|\xi\| \tilde{\triangleright} x = |\xi|^{-1} \langle \xi \rangle \tilde{\triangleleft} \langle \xi \rangle^{-1} = \langle \xi \rangle |\xi|^{-1} = (\langle \xi \rangle \triangleright |\xi|^{-1}) (\langle \xi \rangle \triangleleft |\xi|^{-1}) = v' t',$$

which implies that $t' = \langle \xi \rangle \triangleleft |\xi|^{-1}$, and hence

$$\begin{aligned} \|T(\xi)\| \tilde{\triangleright} x &= \|\xi\| \tilde{\triangleright} x = |\xi|^{-1} \langle \xi \rangle \tilde{\triangleright} \langle \xi \rangle^{-1} = t \langle \xi \rangle^{-1} t'^{-1} \\ &= \langle \xi \rangle \langle \xi \rangle^{-1} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} = (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1}. \end{aligned} \quad \square \quad (29)$$

7 Characters in the category $\mathcal{D}$

Definition 7.1 [9] *The right adjoint action in $\mathcal{D}$ of the algebra D on itself is defined by*




figure 4

Definition 7.2 *The character χ_V of an object V in $\mathcal{D}$ is defined by*




figure 5

Lemma 7.3 *For an object V in $\mathcal{D}$ we have*





figure 6

Proof.



Proposition 7.4 *The character is right adjoint invariant, i.e. for an object V in $\mathcal{D}$*

$$\begin{array}{c} D \quad D \\ | \quad | \\ \textcircled{Ad} \\ | \\ \textcircled{\chi_V} \end{array} = \begin{array}{c} D \quad D \\ | \quad | \\ \textcircled{\chi_V} \quad \textcircled{\epsilon} \end{array}$$

figure 7

Proof.

$$\begin{array}{c} \text{L.H.S.} = \end{array}
 \begin{array}{c} \text{Diagram 1} \end{array}
 =
 \begin{array}{c} \text{Diagram 2} \end{array}
 =
 \begin{array}{c} \text{Diagram 3} \end{array}
 =
 \begin{array}{c} \text{Diagram 4}$$

The proof consists of four diagrams connected by equals signs, showing a sequence of transformations. Each diagram features a central vertical line with various components: a circle labeled θ^{-1} , a circle labeled S , and rectangular blocks. The diagrams are connected by equals signs, indicating a sequence of transformations. The first diagram is labeled "L.H.S. =". The fourth diagram is the final result of the sequence.

The diagram sequence illustrates a proof of an identity in a categorical theory. The components are:

- Octagons:** Represent multiplication and comultiplication. In the first diagram, the left octagon is comultiplication and the right is multiplication. In the second, both are multiplication. In the third, both are comultiplication. In the fourth, both are multiplication. In the fifth, both are comultiplication. In the sixth, both are multiplication.
- Squares:** Represent the Frobenius law. In the first diagram, the top square is comultiplication and the bottom is multiplication. In the second, both are multiplication. In the third, both are comultiplication. In the fourth, both are multiplication. In the fifth, both are comultiplication. In the sixth, both are multiplication.
- Circles:**
 - S : A circle with a vertical line passing through its center, representing the counit.
 - θ^{-1} : A circle with a vertical line passing through its center, representing the unit.
 - ϵ : A circle with a vertical line passing through its center, representing the counit.
- Horizontal Bars:** Represent the multiplication and comultiplication maps.

 The sequence of diagrams is connected by equals signs, showing the following steps:

- Diagram 1: Initial expression with two octagons, two squares, and two S circles.
- Diagram 2: First simplification step, where the octagons are now both multiplication maps.
- Diagram 3: Second simplification step, where the octagons are now both comultiplication maps.
- Diagram 4: Third simplification step, where the octagons are now both multiplication maps.
- Diagram 5: Fourth simplification step, where the octagons are now both comultiplication maps.
- Diagram 6: Final simplification step, where the octagons are now both multiplication maps, and the S circle is replaced by an ϵ circle.

 The final result is labeled "R.H.S" (Right Hand Side) followed by a square symbol $\square$.

Proposition 7.5 *The character of a tensor product of representations is the product of the characters, i.e. for two objects V and W in $\mathcal{D}$*

$$\begin{array}{c} D \\ | \\ \bigcirc \\ \chi_{V \otimes W} \end{array} = \begin{array}{c} D \\ \swarrow \quad \searrow \\ \bigcirc \quad \bigcirc \\ \chi_V \quad \chi_W \end{array}$$


figure 8

Proof.

$$\begin{array}{lcl}
 \text{L.H.S.} = & & \\
 & \begin{array}{c} \text{coev}(V) \\ \text{coev}(W) \\ \text{Diagram 1} \end{array} & = \begin{array}{c} \text{Diagram 2} \end{array} \\
 & \begin{array}{c} \text{Diagram 3} \end{array} & = \begin{array}{c} \text{coev}(V) \quad \text{coev}(W) \\ \text{Diagram 4} \end{array}
 \end{array}$$

The diagrams are as follows:

- Diagram 1:** A vertical line enters from the top, passes through a box labeled θ^{-1} , and then splits into two lines that enter two octagons labeled $\text{coev}(V)$ and $\text{coev}(W)$. The lines from the octagons cross and then exit as a single line at the bottom.
- Diagram 2:** Similar to Diagram 1, but the box labeled θ^{-1} is positioned differently, and the lines from the octagons cross differently.
- Diagram 3:** Similar to Diagram 2, but with a different crossing pattern for the lines from the octagons.
- Diagram 4:** Similar to Diagram 3, but the octagons are labeled $\text{coev}(V)$ and $\text{coev}(W)$ separately, and the box labeled θ^{-1} is positioned differently.



Theorem 7.6 *We have the following formula for the character;*

$$\chi_V(\delta_y \otimes x) = \sum_{\xi \in \text{basis of } V \text{ with } y = \langle \xi \rangle |\xi|^{-1}} \hat{\xi}(\xi \hat{\triangleleft} \langle \xi \rangle^{-1} x \langle \xi \rangle),$$

for $xy = yx$, otherwise $\chi_V(\delta_y \otimes x) = 0$.

Proof. Set $a = \delta_y \otimes x$. To have $\chi_V(a) \neq 0$ we must have $\|a\| = e$, i.e. $y = y \tilde{\triangleleft} x$ which implies that x and y commute. Assuming this, we continue with the diagrammatic definition of the character, starting with

$$\left(\sum_{\xi \in \text{basis of } V} \xi \hat{\triangleleft} \tilde{\tau}(\|\xi\|^L, \|\xi\|)^{-1} \otimes \hat{\xi} \right) \otimes a = \sum_{\xi \in \text{basis of } V} \left(\xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \otimes \hat{\xi} \right) \otimes a.$$

Next we calculate

$$\Psi \left(\xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \otimes \hat{\xi} \right) = \hat{\xi} \hat{\triangleleft} (\langle \xi' \rangle \triangleleft |\hat{\xi}|)^{-1} \otimes \xi' \hat{\triangleleft} |\hat{\xi}| \quad (30)$$

where $\xi' = \xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}$, so

$$\langle \xi' \rangle = \langle \xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \rangle = \langle \xi \bar{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \rangle = \langle \xi \rangle \triangleleft \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}. \quad (31)$$

From a previous calculation we know that $|\hat{\xi}| = \tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1}$, so

$$\begin{aligned} \hat{\xi} \hat{\triangleleft} (\langle \xi' \rangle \triangleleft |\hat{\xi}|)^{-1} &= \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1})^{-1} \\ &= \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \\ \xi' \triangleleft |\hat{\xi}| &= (\xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1}) \hat{\triangleleft} (\tau(\langle \xi \rangle^L, \langle \xi \rangle) |\xi|^{-1}) = \xi \hat{\triangleleft} |\xi|^{-1}, \end{aligned}$$

which gives the next stage in the evaluation of the diagram:

$$\begin{aligned} \sum_{\xi \in \text{basis of } V} \Psi \left(\xi \hat{\triangleleft} \tau(\langle \xi \rangle^L, \langle \xi \rangle)^{-1} \otimes \hat{\xi} \right) \otimes a \\ = \sum_{\xi \in \text{basis of } V} \left(\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes \xi \hat{\triangleleft} |\xi|^{-1} \right) \otimes a. \end{aligned} \quad (32)$$

Now we apply the associator to the last equation to get

$$\begin{aligned} \sum_{\xi \in \text{basis of } V} \Phi \left((\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes \xi \hat{\triangleleft} |\xi|^{-1}) \otimes a \right) \\ = \sum_{\xi \in \text{basis of } V} \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \tilde{\tau}(\|\xi \hat{\triangleleft} |\xi|^{-1}\|^L, \|a\|) \otimes (\xi \hat{\triangleleft} |\xi|^{-1} \otimes a) \\ = \sum_{\xi \in \text{basis of } V} \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \tau(\langle \xi \hat{\triangleleft} |\xi|^{-1} \rangle, e) \otimes (\xi \hat{\triangleleft} |\xi|^{-1} \otimes a) \\ = \sum_{\xi \in \text{basis of } V} \hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes (\xi \hat{\triangleleft} |\xi|^{-1} \otimes (\delta_y \otimes x)) \end{aligned}$$

as $\tau(\langle \xi \hat{\triangleleft} |\xi|^{-1} \rangle, e) = e$. Now apply the action $\hat{\triangleleft}$ to $\xi \hat{\triangleleft} |\xi|^{-1} \otimes (\delta_y \otimes x)$ to get

$$(\xi \hat{\triangleleft} |\xi|^{-1}) \hat{\triangleleft} (\delta_y \otimes x) = \delta_{y, \|\xi \hat{\triangleleft} |\xi|^{-1}\|} (\xi \hat{\triangleleft} |\xi|^{-1}) \hat{\triangleleft} x = \delta_{y, \|\xi\| \|\xi|^{-1}} \xi \hat{\triangleleft} |\xi|^{-1} x, \quad (33)$$

and to get a non-zero answer we must have

$$y = \|\xi\| \tilde{\triangleleft} |\xi|^{-1} = |\xi|^{-1} \langle \xi \rangle \tilde{\triangleleft} |\xi|^{-1} = |\xi| |\xi|^{-1} \langle \xi \rangle |\xi|^{-1} = \langle \xi \rangle |\xi|^{-1}. \quad (34)$$

Thus the character of V is given by

$$\chi_V(\delta_y \otimes x) = \sum_{\xi \in \text{basis of } V \text{ with } y = \langle \xi \rangle |\xi|^{-1}} \text{eval} \left(\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes \theta^{-1}(\xi \hat{\triangleleft} |\xi|^{-1} x) \right).$$

Next

$$\begin{aligned} \theta^{-1}(\xi \hat{\triangleleft} |\xi|^{-1} x) &= (\xi \hat{\triangleleft} |\xi|^{-1} x) \hat{\triangleleft} \|\xi \hat{\triangleleft} |\xi|^{-1} x\|^{-1} \\ &= (\xi \hat{\triangleleft} |\xi|^{-1} x) \hat{\triangleleft} (\|\xi\| \tilde{\triangleleft} |\xi|^{-1} x)^{-1} \\ &= (\xi \hat{\triangleleft} |\xi|^{-1} x) \hat{\triangleleft} (x^{-1} |\xi| |\xi|^{-1} \langle \xi \rangle |\xi|^{-1} x)^{-1} \\ &= \xi \hat{\triangleleft} |\xi|^{-1} x x^{-1} |\xi| \langle \xi \rangle^{-1} x = \xi \hat{\triangleleft} \langle \xi \rangle^{-1} x. \end{aligned}$$

Now we need to calculate $\text{eval}(\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes \xi \hat{\triangleleft} \langle \xi \rangle^{-1} x)$. Start with $\|\xi\| \tilde{\triangleleft} \langle \xi \rangle^{-1} x = \langle \xi \rangle |\xi|^{-1} \tilde{\triangleleft} x = \langle \xi \rangle |\xi|^{-1}$, as we only have nonzero summands for $y = \langle \xi \rangle |\xi|^{-1}$. Then

$$\begin{aligned} &\text{eval}(\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes \xi \hat{\triangleleft} \langle \xi \rangle^{-1} x) \\ &= \text{eval}((\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} \otimes \xi \hat{\triangleleft} \langle \xi \rangle^{-1} x) \hat{\triangleleft} \langle \xi \rangle) \\ &= \text{eval}(\hat{\xi} \hat{\triangleleft} (\langle \xi \rangle \triangleleft |\xi|^{-1})^{-1} (\langle \xi \rangle |\xi|^{-1} \tilde{\triangleright} \langle \xi \rangle) \otimes \xi \hat{\triangleleft} \langle \xi \rangle^{-1} x \langle \xi \rangle). \end{aligned}$$

To find $\langle \xi \rangle |\xi|^{-1} \tilde{\triangleright} \langle \xi \rangle$, first find $\langle \xi \rangle |\xi|^{-1} \tilde{\triangleleft} \langle \xi \rangle = |\xi|^{-1} \langle \xi \rangle$, so

$$\langle \xi \rangle |\xi|^{-1} \tilde{\triangleright} \langle \xi \rangle = (\langle \xi \rangle \triangleright |\xi|^{-1}) (\langle \xi \rangle \triangleleft |\xi|^{-1}) \tilde{\triangleright} \langle \xi \rangle = (\langle \xi \rangle \triangleleft |\xi|^{-1}) \langle \xi \rangle \langle \xi \rangle^{-1} = \langle \xi \rangle \triangleleft |\xi|^{-1}. \quad \square$$

Lemma 7.7 *Let V be an object in $\mathcal{D}$. For $\delta_y \otimes x \in D$ the character of V is given by the following formula, where $y = su^{-1}$ with $s \in M$ and $u \in G$:*

$$\chi_V(\delta_y \otimes x) = \sum_{\xi \in \text{basis of } V_{u^{-1}s}} \hat{\xi}(\xi \hat{\triangleleft} s^{-1} x s) = \chi_{V_{u^{-1}s}}(s^{-1} x s)$$

where $xy = yx$, otherwise $\chi_V(\delta_y \otimes x) = 0$. Here $\chi_{V_{u^{-1}s}}$ is the group representation character of the representation $V_{u^{-1}s}$ of the group $\text{stab}(u^{-1}s)$.

Proof. From theorem 7.6, we know that

$$\chi_V(\delta_y \otimes x) = \sum_{\xi \in \text{basis of } V \text{ with } y = \langle \xi \rangle |\xi|^{-1}} \hat{\xi}(\xi \hat{\Delta} \langle \xi \rangle^{-1} x \langle \xi \rangle),$$

for with $xy = yx$. Set $s = \langle \xi \rangle$ and $u = |\xi|$, so $y = su^{-1}$. We note that $s^{-1}xs$ is in $\text{stab}(u^{-1}s)$, because

$$u^{-1}s\tilde{s}^{-1}xs = s^{-1}x^{-1}su^{-1}ss^{-1}xs = s^{-1}x^{-1}xsu^{-1}s = u^{-1}s.$$

It just remains to note that $\|\xi\| = |\xi|^{-1}\langle \xi \rangle = u^{-1}s$. $\square$

8 Modular Categories

Let $\mathcal{M}$ be a semisimple ribbon category. For objects V and W in $\mathcal{M}$ define $\tilde{S}_{VW} \in \underline{\mathbf{1}}$ by

$$\tilde{S}_{VW} = \begin{array}{c} \text{coev}(V) \quad \text{coev}(W) \\ \begin{array}{c} \text{Diagram showing two octagons representing coevaluation maps for } V \text{ and } W. \text{ The octagons are connected by lines that cross, forming a braid-like structure. Below each octagon is a circle containing } \theta_V^{-1} \text{ and } \theta_W^{-1} \text{ respectively.} \end{array} \end{array}$$

figure 9

There are standard results [1, 13]:

$$\tilde{S}_{VW} = \tilde{S}_{WV} = \tilde{S}_{V^*W^*} = \tilde{S}_{W^*V^*}, \quad \tilde{S}_{V\mathbf{1}} = \dim(V).$$

Here $\dim(V)$ is the trace in $\mathcal{M}$ of the identity map on V .

Definition 8.1 *We call an object U in an abelian category $\mathcal{M}$ simple if, for any V in $\mathcal{M}$, any injection $V \hookrightarrow U$ is either 0 or an isomorphism [1]. A semisimple category is an abelian category whose objects split as a direct sums of simple objects [13].*

Definition 8.2 [1] *A modular category is a semisimple ribbon category $\mathcal{M}$ satisfying the following properties:*

- 1- *There are only a finite number of isomorphism classes of simple objects in $\mathcal{M}$.*
- 2- *Schur's lemma holds, i.e. the morphisms between simple objects are zero unless they are isomorphic, in which case the morphisms are a multiple of the identity.*
- 3- *The matrix $\tilde{S}_{VW}$ with indices in isomorphism classes of simple objects is invertible.*

Definition 8.3 [1] *For a simple object V , the ribbon map on V is a multiple of the identity, and we use Θ_V for the scalar multiple. The numbers $P^\pm$ are defined as the following sums over simple isomorphism classes:*

$$P^\pm = \sum_V \Theta_V^{\pm 1} (\dim(V))^2 ,$$

and the matrices T and C are defined using the Kronecker delta function by

$$T_{VW} = \delta_{VW} \Theta_V , \quad C_{VW} = \delta_{VW^*} .$$

Theorem 8.4 [1] *In a modular category, if we define the matrix S by*

$$S = \frac{\tilde{S}}{\sqrt{P^+ P^-}} ,$$

then we have the following matrix equations:

$$(ST)^3 = \sqrt{\frac{P^+}{P^-}} S^2 , \quad S^2 = C , \quad CT = TC , \quad C^2 = 1 .$$

28

Proof.

$$\begin{aligned}
 L.H.S. &= \text{Diagram 1} = \text{Diagram 2} \\
 &= \text{Diagram 3} = \text{Diagram 4} = R.H.S. \quad \square
 \end{aligned}$$

Lemma 8.7 For V, W indecomposable objects in $\mathcal{D}$, $\text{trace}(\Psi_{V^*W} \circ \Psi_{WV^*}) = \tilde{S}_{VW}$.

Proof.

$$\begin{aligned}
 L.H.S. &= \text{Diagram 1} = \text{Diagram 2}
 \end{aligned}$$

$$\begin{array}{c}
= \\
= \\
= \\
= \\
= \\
=
\end{array}$$

The sequence of diagrams is as follows:

- Diagram 1:** A complex knot with a top loop labeled $\text{coev}(W)$ and $\text{coev}(V)$. It contains circles labeled u , θ_W^{-1} , and $\theta_{V^*}^{-1}$.
- Diagram 2:** A similar knot, but with the $\theta_{V^*}^{-1}$ circle moved to a different position.
- Diagram 3:** The u circle is moved to the bottom of the diagram.
- Diagram 4:** The $\theta_{V^*}^{-1}$ circle is moved to the bottom, next to the u circle.
- Diagram 5:** The θ_W^{-1} circle is moved to the bottom, next to the $\theta_{V^*}^{-1}$ circle.
- Diagram 6:** The strands are rearranged, with the $\text{coev}(V)$ and $\text{coev}(W)$ labels at the top.
- Diagram 7:** The final simplified form, showing two separate loops, each containing a θ circle, with the u circle at the bottom.

= R. H. S. $\square$

Lemma 8.8 For two objects V and W in $\mathcal{D}$,

$$\text{trace}(\Psi_{W \otimes V} \circ \Psi_{V \otimes W}) = \sum_{\substack{\xi \otimes \eta \in \text{basis of } V \otimes W \text{ and} \\ |\xi|^{-1} \langle \xi \rangle \text{ commutes with } |\eta| \langle \eta \rangle^{-1}}} \hat{\eta}(\eta \hat{\Delta} |\eta|^{-1} \langle \xi \rangle^{-1} |\xi| |\eta|) \hat{\xi}(\xi \hat{\Delta} |\eta| \langle \eta \rangle^{-1})$$

Proof. From theorem 6.2, we know that

$$\text{trace}(\Psi_{W \otimes V} \circ \Psi_{V \otimes W}) = \sum_{(\xi \otimes \eta) \in \text{basis of } V \otimes W} \widehat{(\xi \otimes \eta)}(\Psi^2(\xi \otimes \eta)) . \quad (35)$$

From the definition of the ribbon map, we know that $\Psi(\Psi(\xi \otimes \eta)) \hat{\Delta} \|\xi \otimes \eta\| = \xi \hat{\Delta} \|\xi\| \otimes \eta \hat{\Delta} \|\eta\|$, so

$$\begin{aligned} \Psi(\Psi(\xi \otimes \eta)) &= (\xi \hat{\Delta} \|\xi\| \otimes \eta \hat{\Delta} \|\eta\|) \hat{\Delta} \|\xi \otimes \eta\|^{-1} \\ &= (\xi \hat{\Delta} |\xi|^{-1} \langle \xi \rangle \otimes \eta \hat{\Delta} |\eta|^{-1} \langle \eta \rangle) \hat{\Delta} \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta| \\ &= (\xi \hat{\Delta} |\xi|^{-1} \langle \xi \rangle) \hat{\Delta} (\|\eta \hat{\Delta} \|\eta\| \|\tilde{\nu} \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta|\) \otimes \eta \hat{\Delta} |\eta|^{-1} \langle \eta \rangle \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta| \\ &= \xi \hat{\Delta} |\xi|^{-1} \langle \xi \rangle (\|\eta\| \tilde{\nu} \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta|) \otimes \eta \hat{\Delta} |\eta|^{-1} \langle \xi \rangle^{-1} |\xi| |\eta|. \end{aligned}$$

Put $\Psi(\Psi(\xi \otimes \eta)) = \xi' \otimes \eta'$ and $\widehat{\xi \otimes \eta} = \alpha \otimes \beta$, and then from lemma 4.1 we get

$$\widehat{(\xi \otimes \eta)}(\xi' \otimes \eta') = (\alpha \bar{\Delta} \tau(\langle \beta \rangle, \langle \xi' \rangle \cdot \langle \eta' \rangle))(\eta') (\beta \bar{\Delta} \tau(\langle \xi' \rangle, \langle \eta' \rangle)^{-1})(\xi').$$

As $\widehat{\xi \otimes \eta}$ is part of a dual basis, the last expression can only be non-zero if $\|\xi'\| = \|\xi\|$ and $\|\eta'\| = \|\eta\|$. A simple calculation shows that $\|\eta'\| = \|\eta\|$ if and only if $|\xi|^{-1} \langle \xi \rangle$ commutes with $|\eta| \langle \eta \rangle^{-1}$. We use this to find

$$\begin{aligned} \|\eta\| \tilde{\Delta} \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta| &= |\eta|^{-1} |\xi|^{-1} \langle \xi \rangle \langle \eta \rangle |\eta|^{-1} \langle \eta \rangle \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta| \\ &= |\eta|^{-1} \langle \eta \rangle |\eta|^{-1} |\xi|^{-1} \langle \xi \rangle \langle \xi \rangle^{-1} |\xi| |\eta| = |\eta|^{-1} \langle \eta \rangle , \end{aligned}$$

and then

$$\|\eta\| \tilde{\Delta} \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta| = \langle \eta \rangle \langle \eta \rangle^{-1} \langle \xi \rangle^{-1} |\xi| |\eta| \langle \eta \rangle^{-1} = \langle \xi \rangle^{-1} |\xi| |\eta| \langle \eta \rangle^{-1} .$$

Now using the formula for $\widehat{\xi \otimes \eta} = \alpha \otimes \beta$ from lemma 4.1 gives the result. $\square$

Lemma 8.9 *Let V and W be objects in $\mathcal{D}$. Then in terms of group characters:*

$$\text{trace}(\Psi^2_{V \otimes W}) = \sum_{\substack{u,v \in G, s,t \in M \text{ and} \\ su \text{ commutes with } vt}} \chi_{W_{us}}(s^{-1}t^{-1}v^{-1}s) \chi_{V_{vt}}(u^{-1}s^{-1}).$$

Proof. This is more or less immediate from lemma 8.8. Put $\|\eta\| = u^{-1}s$ and $\|\xi\| = v^{-1}t$ and sum over basis elements of constant degree first. $\square$

9 An example of a modular category

Using the order of the indecomposable objects in table (5), we get T to be a diagonal 32×32 matrix whose diagonal entries are taken from the table. As every indecomposable object in our example is self-dual, the matrix C is the 32×32 identity matrix.

To find S , we calculate the trace of the double braiding $\text{trace}(\Psi_{VW} \circ \Psi_{WV})$. We do this using the result from 8.8, and split into different cases for the objects V and W , and move the points the characters are evaluated at to the base points for each orbit using 2.3. For example:

Case (1) $\otimes$ Case (1): (i.e. the orbit of W is $\{e\}$ and the orbit of V is $\{e\}$)

$$\text{trace}(\Psi^2) = \chi_{W_e}(e)\chi_{V_e}(e).$$

Case (2) $\otimes$ Case (5): (i.e. the orbit of W is $\{a^3\}$ and the orbit of V is $\{b, ba^2, ba^4\}$)

$$\text{trace}(\Psi^2) = \left(\chi_{W_{a^3}}(ba^2) + \chi_{W_{a^3}}(ba^4) + \chi_{W_{a^3}}(b) \right) \chi_{V_b}(a^3).$$

Case (5) $\otimes$ Case (3): (i.e. the orbit of W is $\{b, ba^2, ba^4\}$ and the orbit of V is $\{a^2, a^4\}$)

$$\text{trace}(\Psi^2) = 0.$$

Case (6) $\otimes$ Case (5): (i.e. the orbit of W is $\{ba, ba^3, ba^5\}$ and the orbit of V is $\{b, ba^2, ba^4\}$)

$$\text{trace}(\Psi^2) = 3 \left(\chi_{W_{ba}}(ba^4) \chi_{V_b}(ba^3) \right).$$

From these cases we get S to be the following 32×32 symmetric matrix:

[illegible]

Now it is possible to check that the matrices S , T and C satisfy the following relations:

$$S^2 = \frac{1}{12}(ST)^3, \quad CS = SC, \quad CT = TC.$$

10 An equivalence of tensor categories

In this section we will generalize some results of [3] which considered group doublecross products, i.e. a group X factoring into two subgroups G and M .

Definition 10.1 [3] *For the doublecross product group $X = GM$ there is a quantum double $D(X) = k(X) \bowtie kX$ which has the following operations*

$$(\delta_y \otimes x)(\delta_{y'} \otimes x') = \delta_{x^{-1}yx, y'}(\delta_y \otimes xx'), \quad \Delta(\delta_y \otimes x) = \sum_{ab=y} \delta_a \otimes x \otimes \delta_b \otimes x$$

$$1 = \sum_y \delta_y \otimes e, \quad \epsilon(\delta_y \otimes x) = \delta_{y,e}, \quad S(\delta_y \otimes x) = \delta_{x^{-1}y^{-1}x} \otimes x^{-1},$$

$$(\delta_y \otimes x)^* = \delta_{x^{-1}yx} \otimes x^{-1}, \quad R = \sum_{x,z} \delta_x \otimes e \otimes \delta_z \otimes x.$$

The representations of $D(X)$ are given by X -graded left kX -modules. The kX action will be denoted by $\triangleright$ and the grading by $\|\cdot\|$. The grading and X action are related by

$$\|x \triangleright \xi\| = x \|\xi\| x^{-1} \quad x \in X, \quad \xi \in V, \quad (36)$$

and the action of $(\delta_y \otimes x) \in D(X)$ is given by

$$(\delta_y \otimes x) \triangleright \xi = \delta_{y, \|x \triangleright \xi\|} x \triangleright \xi. \quad (37)$$

Proposition 10.2 *There is a functor χ from $\mathcal{D}$ to the category of representations of $D(X)$ given by the following: As vector spaces, $\chi(V)$ is the same as V , and χ is the identity map. The X -grading $\|\cdot\|$ on $\chi(V)$ and the action of $us \in kX$ are defined by*

$$\begin{aligned} \|\chi(\eta)\| &= \langle \eta \rangle^{-1} |\eta| \quad \text{for } \eta \in V, \\ us \triangleright \chi(\eta) &= \chi\left(\left((s \triangleleft |\eta|^{-1}) \triangleright \eta\right) \bar{\triangleleft} u^{-1}\right), \quad s \in M \quad u \in G. \end{aligned}$$

A morphism $\phi : V \rightarrow W$ in $\mathcal{D}$ is sent to the morphism $\chi(\phi) : \chi(V) \rightarrow \chi(W)$ defined by $\chi(\phi)(\chi(\xi)) = \chi(\phi(\xi))$.

Proof. First we show that $\triangleright$ is an action, i.e. $vt\triangleright(us\triangleright\chi(\eta)) = vtus\triangleright\chi(\eta)$ for all $s, t \in M$ and $u, v \in G$. Note that

$$\begin{aligned} vt\triangleright(us\triangleright\chi(\eta)) &= vt\triangleright\chi((s\triangleleft|\eta|^{-1})\bar{\triangleright}\eta)\bar{\triangleleft}u^{-1}) \\ &= \chi(((t\triangleleft|\bar{\eta}|^{-1})\bar{\triangleright}\bar{\eta})\bar{\triangleleft}v^{-1}), \end{aligned}$$

where $\bar{\eta} = ((s\triangleleft|\eta|^{-1})\bar{\triangleright}\eta)\bar{\triangleleft}u^{-1}$. On the other hand we have

$$vtus = v(t\triangleright u)\tau(t\triangleleft u, s)((t\triangleleft u) \cdot s),$$

where $v(t\triangleright u)\tau(t\triangleleft u, s) \in G$ and $(t\triangleleft u) \cdot s \in M$, so

$$vtus\triangleright\chi(\eta) = \chi\left(\left(\left(\left((t\triangleleft u) \cdot s\right)\triangleleft|\eta|^{-1}\right)\bar{\triangleright}\eta\right)\bar{\triangleleft}\tau(t\triangleleft u, s)^{-1}(t\triangleright u)^{-1}v^{-1}\right) \right).$$

We need to show that

$$\begin{aligned} (t\triangleleft|\bar{\eta}|^{-1})\bar{\triangleright}\bar{\eta} &= \left(\left(\left((t\triangleleft u) \cdot s\right)\triangleleft|\eta|^{-1}\right)\bar{\triangleright}\eta\right)\bar{\triangleleft}\tau(t\triangleleft u, s)^{-1}(t\triangleright u)^{-1} \right. \\ &= \left(\left((t\triangleleft u(s\triangleright|\eta|^{-1})) \cdot (s\triangleleft|\eta|^{-1})\right)\bar{\triangleright}\eta\right)\bar{\triangleleft}\tau(t\triangleleft u, s)^{-1}(t\triangleright u)^{-1} \end{aligned} \quad (38)$$

Put $\bar{s} = s\triangleleft|\eta|^{-1}$ and $\eta' = \bar{s}\bar{\triangleright}\eta$ which give $\bar{\eta} = \eta'\bar{\triangleleft}u^{-1}$. Then using the connections between the gradings and actions,

$$|\bar{\eta}| = |\eta'\bar{\triangleleft}u^{-1}| = (\langle\eta'\rangle\triangleright u^{-1})^{-1}|\eta'|u^{-1}.$$

Putting $\bar{t} = t\triangleleft u|\eta'|^{-1}$, the left hand side of (38) will become

$$\begin{aligned} (t\triangleleft|\bar{\eta}|^{-1})\bar{\triangleright}\bar{\eta} &= \left(t\triangleleft u|\eta'|^{-1}(\langle\eta'\rangle\triangleright u^{-1})\right)\bar{\triangleright}(\eta'\bar{\triangleleft}u^{-1}) \\ &= \left(\bar{t}\triangleleft(\langle\eta'\rangle\triangleright u^{-1})\right)\bar{\triangleright}(\eta'\bar{\triangleleft}u^{-1}) \\ &= (\bar{t}\bar{\triangleright}\eta')\bar{\triangleleft}((\bar{t}\triangleleft|\eta'|)\triangleright u^{-1}). \end{aligned}$$

Now, from (38) and the fact that $(t\triangleright u)^{-1} = (\bar{t}\triangleleft|\eta'|)\triangleright u^{-1}$, we only need to show that

$$\bar{t}\bar{\triangleright}\eta' = \left(\left((t\triangleleft u(s\triangleright|\eta|^{-1})) \cdot (s\triangleleft|\eta|^{-1})\right)\bar{\triangleright}\eta\right)\bar{\triangleleft}\tau(t\triangleleft u, s)^{-1}. \quad (39)$$

From the formula for the composition of the M ‘action’ the right hand side of (39) becomes $\bar{p} \bar{\triangleright} (\bar{s} \bar{\triangleright} \eta) = \bar{p} \bar{\triangleright} \eta'$, where $\bar{p}' = t \triangleleft u(s \triangleright |\eta|^{-1})$ and $\bar{p} = \bar{p}' \triangleleft \tau(\bar{s}, \langle \eta \rangle) \tau(\langle \bar{s} \bar{\triangleright} \eta \rangle, \bar{s} \triangleleft |\eta|)^{-1}$. We have used the fact that $\tau(t \triangleleft u, s) = \tau(\bar{p}' \triangleleft (\bar{s} \triangleright |\eta|), \bar{s} \triangleleft |\eta|)$. Now we just have to prove that $\bar{p} = \bar{t}$. Because $\tau(\bar{s}, \langle \eta \rangle)^{-1} (\bar{s} \triangleright |\eta|) = \tau(\langle \bar{s} \bar{\triangleright} \eta \rangle, \bar{s} \triangleleft |\eta|)^{-1} |\bar{s} \bar{\triangleright} \eta|$ and knowing that $(\bar{s} \triangleright |\eta|) = (s \triangleright |\eta|^{-1})^{-1}$, we can write $\bar{p}$ as follows

$$\begin{aligned} \bar{p} &= \bar{p}' \triangleleft (\bar{s} \triangleright |\eta|) |\bar{s} \triangleright \eta|^{-1} \\ &= t \triangleleft u(s \triangleright |\eta|^{-1}) (s \triangleright |\eta|^{-1})^{-1} |\eta'|^{-1} \\ &= t \triangleleft u |\eta'|^{-1} = \bar{t}. \end{aligned}$$

Next we show that $\|us \dot{\chi}(\eta)\| = us \|\chi(\eta)\| (us)^{-1}$ where $u \in G$ and $s \in M$.

$$\begin{aligned} \|x \dot{\chi}(\eta)\| &= \|\chi((s \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta) \bar{\triangleleft} u^{-1})\| \\ &= \langle \eta' \bar{\triangleleft} u^{-1} \rangle^{-1} |\eta' \bar{\triangleleft} u^{-1}| = u \langle \eta' \rangle^{-1} |\eta'| u^{-1} \\ &= u \langle \bar{s} \bar{\triangleright} \eta \rangle^{-1} |\bar{s} \bar{\triangleright} \eta| u^{-1} = u (\bar{s} \triangleleft |\eta|) \langle \eta \rangle^{-1} |\eta| (\bar{s} \triangleleft |\eta|)^{-1} u^{-1} \\ &= us \langle \eta \rangle^{-1} |\eta| s^{-1} u^{-1}. \quad \square \end{aligned}$$

Theorem 10.3 *The functor χ is invertible.*

Proof. We have already proved in the previous proposition that the X -grading $\|\cdot\|$ and the action $\dot{\triangleright}$ give a representation of $D(X)$, so we only need to show that χ is a 1-1 correspondence, which we do by giving its inverse χ^{-1} as the following: Let W be a representation of $D(X)$, with kX action $\dot{\triangleright}$ and X -grading $\|\cdot\|$. Define a D representation as follows: $\chi^{-1}(W)$ will be the same as W as a vector space. There will be G and M gradings given by the factorization

$$\|\xi\|^{-1} = |\chi^{-1}(\xi)|^{-1} \langle \chi^{-1}(\xi) \rangle, \quad \xi \in W, \quad \langle \chi^{-1}(\xi) \rangle \in M, \quad |\chi^{-1}(\xi)| \in G.$$

The action of $s \in M$ and $u \in G$ are given by

$$s\bar{\triangleright}\chi^{-1}(\xi) = \chi^{-1}((s\triangleleft|\chi^{-1}(\xi)|)\dot{\triangleright}\xi) , \quad \chi^{-1}(\xi)\bar{\triangleleft}u = \chi^{-1}(u^{-1}\dot{\triangleright}\xi)$$

Checking the rest is left to the reader. $\square$

Proposition 10.4 *For $\delta_y \otimes x \in \mathcal{D}$, $\chi(\xi\hat{\triangleleft}(\delta_y \otimes x)) = \delta_{y,\|\xi\|} x^{-1}\dot{\triangleright}\chi(\xi)$.*

Proof. Starting with the left hand side,

$$\chi(\xi\hat{\triangleleft}(\delta_y \otimes x)) = \chi(\delta_{y,\|\xi\|} \xi\hat{\triangleleft}x) = \delta_{y,\|\xi\|} \chi(\xi\hat{\triangleleft}x).$$

Putting $x = us$ for $u \in G$ and $s \in M$,

$$\xi\hat{\triangleleft}x = \xi\hat{\triangleleft}us = (\xi\bar{\triangleleft}u)\hat{\triangleleft}s = ((s^L\triangleleft u^{-1}|\xi|^{-1})\bar{\triangleright}\xi)\bar{\triangleleft}(s^L\triangleright u^{-1})^{-1}\tau(s^L, s).$$

Now put $\bar{u} = \tau(s^L, s)^{-1}(s^L\triangleright u^{-1})$ and $\bar{s} = s^L\triangleleft u^{-1}$. Then

$$\begin{aligned} \chi(\xi\hat{\triangleleft}(\delta_y \otimes x)) &= \delta_{y,\|\xi\|} \chi\left((\bar{s}\triangleleft|\xi|^{-1})\bar{\triangleright}\xi\right)\bar{\triangleleft}\bar{u}^{-1}) \\ &= \delta_{y,\|\xi\|} \bar{u}\bar{s}\dot{\triangleright}\chi(\xi) \\ &= \delta_{y,\|\xi\|} \tau(s^L, s)^{-1}(s^L\triangleright u^{-1})(s^L\triangleleft u^{-1})\dot{\triangleright}\chi(\xi) \\ &= \delta_{y,\|\xi\|} s^{-1}s^{L-1}s^L u^{-1}\dot{\triangleright}\chi(\xi) \\ &= \delta_{y,\|\xi\|} (us)^{-1}\dot{\triangleright}\chi(\xi) = \delta_{y,\|\xi\|} x^{-1}\dot{\triangleright}\chi(\xi). \quad \square \end{aligned}$$

Proposition 10.5 *Define a map $\psi : D \longrightarrow D(X)$ by $\psi(\delta_y \otimes x) = \delta_{x^{-1}yx} \otimes x^{-1}$. Then ψ satisfies the equation $\chi(\xi\hat{\triangleleft}(\delta_y \otimes x)) = \psi(\delta_y \otimes x)\dot{\triangleright}\chi(\xi)$.*

Proof. Use the previous proposition. $\square$

The reader will recall that D is in general a non-trivially associated algebra (i.e. it is only associative in the category $\mathcal{D}$ with its non-trivial associator). Thus, in general, it

can not be isomorphic to $D(X)$, which really is associative. In general, ψ can not be an algebra map.

Proposition 10.6 *For a and b elements of the algebra D in the category $\mathcal{D}$,*

$$\psi(b)\psi(a) = \psi(ab) \left(\sum_{y \in Y} \delta_y \otimes \tau(\langle a \rangle, \langle b \rangle)^{-1} \right).$$

Proof. by 10.5 we have

$$\begin{aligned} \chi((\xi \hat{\triangleleft} a) \hat{\triangleleft} b) &= \psi(b) \dot{\triangleright} \chi(\xi \hat{\triangleleft} a) \\ &= \psi(b) \dot{\triangleright} (\psi(a) \dot{\triangleright} \chi(\xi)) \\ &= \psi(b) \psi(a) \dot{\triangleright} \chi(\xi). \end{aligned}$$

But also, where $f = \sum_y \delta_y \otimes \tau(\langle a \rangle, \langle b \rangle)$,

$$\begin{aligned} \chi((\xi \hat{\triangleleft} a) \hat{\triangleleft} b) &= \chi\left((\xi \hat{\triangleleft} \tilde{\tau}(\|a\|, \|b\|)) \hat{\triangleleft} ab\right) \\ &= \psi(ab) \dot{\triangleright} \chi(\xi \hat{\triangleleft} \tilde{\tau}(\|a\|, \|b\|)) = \psi(ab) \dot{\triangleright} \chi(\xi \hat{\triangleleft} \tilde{\tau}(\langle a \rangle, \langle b \rangle)) \\ &= \psi(ab) \psi(f) \dot{\triangleright} \chi(\xi). \quad \square \end{aligned}$$

Definition 10.7 *Let V and W be objects of $\mathcal{D}$. The map $c : \chi(V) \otimes \chi(W) \longrightarrow \chi(V \otimes W)$*

is defined by:

$$c(\chi(\eta) \otimes \chi(\xi)) = \chi\left(\left((\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta\right) \otimes \xi\right).$$

Proposition 10.8 *The map c , defined above, is a $D(X)$ module map, i.e.*

$$\begin{aligned} \|c(\chi(\eta) \otimes \chi(\xi))\| &= \|\chi(\eta) \otimes \chi(\xi)\|, \\ x \dot{\triangleright} c(\chi(\eta) \otimes \chi(\xi)) &= c\left(x \dot{\triangleright} (\chi(\eta) \otimes \chi(\xi))\right) \quad \forall x \in X. \end{aligned}$$

Proof. We will begin with the grading first. It is known that

$$\|\chi(\eta) \otimes \chi(\xi)\| = \|\chi(\eta)\| \|\chi(\xi)\| = \langle \eta \rangle^{-1} |\eta| \langle \xi \rangle^{-1} |\xi|.$$

But on the other hand we know, from the definition of c , that

$$\begin{aligned} \|c(\chi(\eta) \otimes \chi(\xi))\| &= \|\chi((\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta \otimes \xi)\| \\ &= \langle (\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta \otimes \xi \rangle^{-1} |(\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta \otimes \xi| \\ &= \langle \xi \rangle^{-1} \langle \bar{\eta} \rangle^{-1} |\bar{\eta}| |\xi| \\ &= \langle \xi \rangle^{-1} \langle \bar{s} \bar{\triangleright} \eta \rangle^{-1} |\bar{s} \bar{\triangleright} \eta| |\xi| \\ &= \langle \xi \rangle^{-1} (\bar{s} \triangleleft |\eta|) \langle \eta \rangle^{-1} |\eta| (\bar{s} \triangleleft |\eta|)^{-1} |\xi| \\ &= \langle \eta \rangle^{-1} |\eta| \langle \xi \rangle^{-1} |\xi|, \end{aligned}$$

where $\bar{s} = \langle \xi \rangle \triangleleft |\eta|^{-1}$ and $\bar{\eta} = (\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta = \bar{s} \bar{\triangleright} \eta$, which gives the result.

For the G action, we know from the definitions that

$$\begin{aligned} u \dot{\triangleright} (\chi(\eta) \otimes \chi(\xi)) &= \chi(\eta \bar{\triangleleft} u^{-1}) \otimes \chi(\xi \bar{\triangleleft} u^{-1}), \\ c(u \dot{\triangleright} (\chi(\eta) \otimes \chi(\xi))) &= \chi\left(\left((\langle \xi \bar{\triangleleft} u^{-1} \rangle \triangleleft |\eta \bar{\triangleleft} u^{-1}|^{-1}) \bar{\triangleright} (\eta \bar{\triangleleft} u^{-1})\right) \otimes (\xi \bar{\triangleleft} u^{-1})\right). \end{aligned}$$

By using the properties of the G and M gradings,

$$\begin{aligned} \langle \xi \bar{\triangleleft} u^{-1} \rangle \triangleleft |\eta \bar{\triangleleft} u^{-1}|^{-1} &= (\langle \xi \rangle \triangleleft u^{-1}) \triangleleft |\eta|^{-1} (\langle \eta \rangle \triangleright u^{-1}) \\ &= \langle \xi \rangle \triangleleft |\eta|^{-1} (\langle \eta \rangle \triangleright u^{-1}) \\ (\langle \xi \bar{\triangleleft} u^{-1} \rangle \triangleleft |\eta \bar{\triangleleft} u^{-1}|^{-1}) \bar{\triangleright} (\eta \bar{\triangleleft} u^{-1}) &= ((\langle \xi \rangle \triangleleft |\eta|^{-1}) \triangleleft (\langle \eta \rangle \triangleright u^{-1})) \bar{\triangleright} (\eta \bar{\triangleleft} u^{-1}) \\ &= ((\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta) \bar{\triangleleft} \left((\langle \xi \rangle \triangleleft |\eta|^{-1}) \triangleleft |\eta|\right) \triangleright u^{-1} \\ &= ((\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta) \bar{\triangleleft} (\langle \xi \rangle \triangleright u^{-1}). \end{aligned}$$

Now we can write

$$c(u \dot{\triangleright} (\chi(\eta) \otimes \chi(\xi))) = \chi\left(\left((\langle \xi \rangle \triangleleft |\eta|^{-1}) \bar{\triangleright} \eta\right) \bar{\triangleleft} (\langle \xi \rangle \triangleright u^{-1}) \otimes (\xi \bar{\triangleleft} u^{-1})\right). \quad (40)$$

On the other hand,

$$\begin{aligned} u\dot{\triangleright}c(\chi(\eta) \otimes \chi(\xi)) &= u\dot{\triangleright}\chi\left(\left(\langle \xi \rangle \triangleleft |\eta|^{-1}\right) \bar{\triangleright} \eta\right) \otimes \xi \\ &= \chi\left(\left(\left(\langle \xi \rangle \triangleleft |\eta|^{-1}\right) \bar{\triangleright} \eta\right) \otimes \xi\right) \bar{\triangleleft} u^{-1}, \end{aligned}$$

which gives the same as (40).

Now we show that c preserves the M action. For $s \in M$,

$$\begin{aligned} s\dot{\triangleright}(\chi(\eta) \otimes \chi(\xi)) &= \chi((s\triangleleft |\eta|^{-1}) \bar{\triangleright} \eta) \otimes \chi((s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi) \\ c\left(s\dot{\triangleright}(\chi(\eta) \otimes \chi(\xi))\right) &= \chi\left(\left(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft |(s\triangleleft |\eta|^{-1}) \bar{\triangleright} \eta|^{-1}\right) \bar{\triangleright} ((s\triangleleft |\eta|^{-1}) \bar{\triangleright} \eta) \right. \\ &\quad \left. \otimes ((s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi)\right). \end{aligned}$$

Using the ‘action’ property for $\bar{\triangleright}$, we get

$$\left(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft |(s\triangleleft |\eta|^{-1}) \bar{\triangleright} \eta|^{-1}\right) \bar{\triangleright} ((s\triangleleft |\eta|^{-1}) \bar{\triangleright} \eta) = ((p' \cdot \bar{t}) \bar{\triangleright} \eta) \bar{\triangleleft} \tau(p' \triangleleft (\bar{t} \triangleright |\eta|), \bar{t} \triangleleft |\eta|)^{-1},$$

where $\bar{t} = s\triangleleft |\eta|^{-1}$ and

$$p' = \langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft |\bar{t} \bar{\triangleright} \eta|^{-1} \tau(\langle \bar{t} \bar{\triangleright} \eta \rangle, \bar{t} \triangleleft |\eta|) \tau(\bar{t}, \langle \eta \rangle)^{-1}.$$

But using the connections between the grading and the actions, we know that $|\bar{t} \bar{\triangleright} \eta|^{-1} =$

$(\bar{t} \triangleright |\eta|)^{-1} \tau(\bar{t}, \langle \eta \rangle) \tau(\langle \bar{t} \bar{\triangleright} \eta \rangle, \bar{t} \triangleleft |\eta|)^{-1}$, so

$$\begin{aligned} p' &= \langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft (\bar{t} \triangleright |\eta|)^{-1} \\ &= \langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft ((s\triangleleft |\eta|^{-1}) \triangleright |\eta|)^{-1} \\ &= \langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft (s \triangleright |\eta|^{-1}). \end{aligned}$$

Substituting in the equation above gives

$$\begin{aligned} &\left(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft |(s\triangleleft |\eta|^{-1}) \bar{\triangleright} \eta|^{-1}\right) \bar{\triangleright} ((s\triangleleft |\eta|^{-1}) \bar{\triangleright} \eta) \\ &= \left(\left(\left(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \triangleleft (s \triangleright |\eta|^{-1})\right) \cdot (s\triangleleft |\eta|^{-1})\right) \bar{\triangleright} \eta\right) \bar{\triangleleft} \tau(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle, s)^{-1} \\ &= \left(\left(\left(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle \cdot s\right) \triangleleft |\eta|^{-1}\right) \bar{\triangleright} \eta\right) \bar{\triangleleft} \tau(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle, s)^{-1} \\ &= \left(\left(\left((s\triangleleft |\xi|^{-1}) \cdot \langle \xi \rangle\right) \triangleleft |\eta|^{-1}\right) \bar{\triangleright} \eta\right) \bar{\triangleleft} \tau(\langle (s\triangleleft |\xi|^{-1}) \bar{\triangleright} \xi \rangle, s)^{-1}. \end{aligned}$$

On the other hand, we know that

$$\begin{aligned} s\dot{\triangleright}c(\chi(\eta) \otimes \chi(\xi)) &= s\dot{\triangleright}\chi\left(\left((\langle\xi\rangle\triangleleft|\eta|^{-1})\bar{\triangleright}\eta\right) \otimes \xi\right) \\ &= s\dot{\triangleright}\chi(\bar{\eta} \otimes \xi) = \chi\left((s\triangleleft|\bar{\eta} \otimes \xi|^{-1})\bar{\triangleright}(\bar{\eta} \otimes \xi)\right), \end{aligned}$$

where $\bar{\eta} = (\langle\xi\rangle\triangleleft|\eta|^{-1})\bar{\triangleright}\eta$. Next we calculate

$$\begin{aligned} |\bar{\eta} \otimes \xi| &= \tau(\langle\bar{\eta}\rangle, \langle\xi\rangle)^{-1}|\bar{\eta}||\xi|, \\ s\triangleleft|\bar{\eta} \otimes \xi|^{-1} &= s\triangleleft|\xi|^{-1}|\bar{\eta}|^{-1}\tau(\langle\bar{\eta}\rangle, \langle\xi\rangle). \end{aligned}$$

If we put $\bar{s} = s\triangleleft|\xi|^{-1}|\bar{\eta}|^{-1}$, then

$$\begin{aligned} (s\triangleleft|\bar{\eta} \otimes \xi|^{-1})\bar{\triangleright}(\bar{\eta} \otimes \xi) &= (\bar{s}\triangleleft\tau(\langle\bar{\eta}\rangle, \langle\xi\rangle))\bar{\triangleright}(\bar{\eta} \otimes \xi) \\ &= (\bar{s}\bar{\triangleright}\bar{\eta})\bar{\triangleleft}\tau(\bar{s}\triangleleft|\bar{\eta}|, \langle\xi\rangle)\tau(\langle(\bar{s}\triangleleft|\bar{\eta}|)\bar{\triangleright}\xi\rangle, \bar{s}\triangleleft|\bar{\eta}||\xi|)^{-1} \\ &\quad \otimes (\bar{s}\triangleleft|\bar{\eta}|)\bar{\triangleright}\xi \\ &= (\bar{s}\bar{\triangleright}\bar{\eta})\bar{\triangleleft}\tau(s\triangleleft|\xi|^{-1}, \langle\xi\rangle)\tau(\langle(s\triangleleft|\xi|^{-1})\bar{\triangleright}\xi\rangle, s)^{-1} \\ &\quad \otimes (s\triangleleft|\xi|^{-1})\bar{\triangleright}\xi. \end{aligned}$$

Using the ‘action’ property again,

$$\begin{aligned} \bar{s}\bar{\triangleright}\bar{\eta} &= (s\triangleleft|\xi|^{-1}|\bar{\eta}|^{-1})\bar{\triangleright}((\langle\xi\rangle\triangleleft|\eta|^{-1})\bar{\triangleright}\eta) \\ &= \left((q' \cdot (\langle\xi\rangle\triangleleft|\eta|^{-1}))\bar{\triangleright}\eta\right)\bar{\triangleleft}\tau(q'\triangleleft((\langle\xi\rangle\triangleleft|\eta|^{-1})\triangleright|\eta|), \langle\xi\rangle)^{-1} \\ &= \left((q' \cdot (\langle\xi\rangle\triangleleft|\eta|^{-1}))\bar{\triangleright}\eta\right)\bar{\triangleleft}\tau(q'\triangleleft(\langle\xi\rangle\triangleright|\eta|^{-1})^{-1}, \langle\xi\rangle)^{-1}, \end{aligned}$$

where

$$\begin{aligned} q' &= (s\triangleleft|\xi|^{-1}|\bar{\eta}|^{-1})\triangleleft\tau(\langle(\langle\xi\rangle\triangleleft|\eta|^{-1})\triangleright\eta\rangle, \langle\xi\rangle)\tau(\langle\xi\rangle\triangleleft|\eta|^{-1}, \langle\eta\rangle)^{-1} \\ &= (s\triangleleft|\xi|^{-1})\triangleleft(\langle\xi\rangle\triangleright|\eta|^{-1}), \end{aligned}$$

as

$$|\bar{\eta}|^{-1} = ((\langle\xi\rangle\triangleleft|\eta|^{-1})\triangleright|\eta|)^{-1}\tau(\langle\xi\rangle\triangleleft|\eta|^{-1}, \langle\eta\rangle)\tau(\langle(\langle\xi\rangle\triangleleft|\eta|^{-1})\triangleright\eta\rangle, \langle\xi\rangle)^{-1}.$$

Hence substituting with the value of q' we get

$$\begin{aligned}\bar{s}\bar{\triangleright}\bar{\eta} &= \left(\left(((s\triangleleft|\xi|^{-1})\triangleleft(\langle\xi\rangle\triangleright|\eta|^{-1})) \cdot (\langle\xi\rangle\triangleleft|\eta|^{-1}) \right) \bar{\triangleright}\eta \right) \bar{\triangleleft} \tau((s\triangleleft|\xi|^{-1}), \langle\xi\rangle)^{-1} \\ &= \left(\left(((s\triangleleft|\xi|^{-1}) \cdot \langle\xi\rangle) \triangleleft|\eta|^{-1} \right) \bar{\triangleright}\eta \right) \bar{\triangleleft} \tau(s\triangleleft|\xi|^{-1}, \langle\xi\rangle)^{-1},\end{aligned}$$

giving the required result

$$\begin{aligned}(\bar{s}\bar{\triangleright}\bar{\eta})\bar{\triangleleft} \tau(s\triangleleft|\xi|^{-1}, \langle\xi\rangle) \tau(\langle(s\triangleleft|\xi|^{-1})\bar{\triangleright}\xi\rangle, s)^{-1} \\ = \left(\left(((s\triangleleft|\xi|^{-1}) \cdot \langle\xi\rangle) \triangleleft|\eta|^{-1} \right) \bar{\triangleright}\eta \right) \bar{\triangleleft} \tau(\langle(s\triangleleft|\xi|^{-1})\bar{\triangleright}\xi\rangle, s)^{-1} .\square\end{aligned}$$

References

- [1] BAKALOV B. & KIRILLOV A., *Lectures on Tensor Categories and Modular Functors*. American Mathematical Society, 2000.
- [2] BEGGS E. J., *Making non-trivially associated tensor categories from left coset representatives*. *Journal of Pure and Applied Algebra*, vol 177, 5 - 41, 2003.
- [3] BEGGS E. J., GOULD J. D. AND MAJID S., *Finite group factorizations and braiding*. *J. Algebra*, vol 181 no. 1, 112 - 151, 1996.
- [4] BEGGS E. J. AND MAJID S., *Quasitriangular and differential structures on bi-crossproduct Hopf algebras*. *J. Algebra*, vol 219 no. 2, 682 - 727, 1999.
- [5] GROVE L. C., *Groups and Characters*. Wiley-Interscience, 1997.
- [6] GUREVICH D. I. AND MAJID S., *Braided groups of Hopf algebras obtained by twisting*. *Pacific J. Math*, vol 162, 27 - 44, 1994.

- [7] MAJID S., *Physics for algebraists: Non-commutative and non-cocommutative Hopf algebras by bicrossproduct construction.* *J. Algebra*, vol 130, 17 - 64, 1990.
From PhD Thesis, Harvard, 1988.
- [8] MAJID S., *The quantum double as quantum mechanics.* *J. Geom. Phys.*, vol 13, 169 - 202, 1994.
- [9] MAJID S., *Foundations of Quantum Group Theory.* Cambridge University Press, 1995.
- [10] TAKEUCHI M. , *Matched pairs of groups and bismash products of Hopf algebras.* *Commun. Alg.*, vol. 9, no. 8, 841-882 , 1981.
- [11] TAKEUCHI M. , *Modular Categories and Hopf Algebras.* *Journal of Algebra*, vol. 243, no. 2, 631-643, 2001.
- [12] THOMAS A. D. & WOOD G. V., *Group Tables.* Shiva Publishing Ltd., 1980.
- [13] TURAEV V. & WENZL H., *Semisimple and modular categories from link invariants.* *Math. Ann.*, vol. 309, no. 3, 411-461, 1997.